

AD-A254 496



## INTATION PAGE

ACT

Form Approved  
OMB No. 0704-0188

total to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, reviewing the collection of information, Send comments regarding this burden estimate or any other aspect of this burden, to Washington Headquarters Service, Directorate for Information Operations and Reports, 1215 Jefferson Avenue, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE		3. REPORT TYPE AND DATES COVERED FINAL 01 Dec 88 TO 31 May 90	
4. TITLE AND SUBTITLE CARBOXYLESTERASES OF THE TESTES: ROLE IN ACTIVATION OF TOXICANTS				5. FUNDING NUMBERS GR - AFOSR-89-0190 PE - 61102f PR - 2312 TA - A5	
6. AUTHOR(S) Dr Alexandra Ventura					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) American College of Toxicology 9650 Rockville Pike Bethesda, MD 20814					
8. PERFORMING ORGANIZATION REPORT NUMBER AFOSR-TR-92-0800				9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) AFOSR/NL Building 410 Bolling AFB DC 20332/6448	
10. SPONSORING / MONITORING AGENCY REPORT NUMBER				11. SUPPLEMENTARY NOTES	
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited				12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Organ specific distribution of carboxylesterases (Western blotting) was determined to be liver ] lung = testes = fat ] pancreas ] kidney. Carboxylesterase distribution among cell types of the testes was examined by in situ hybridization techniques. Results were inconclusive, as both the probe and the control hybridized to tissues macromolecules. More refinement of this techniques should provide better results. Other accomplishments include examination of the down-regulation of carboxylesterase levels by glucocorticoids. Apparently esterase levels are most dramatically down-regulated (approximately 6-fold) by dexamethasone phosphate (60 mg/kg x 5 days, i.p.) in the testes compared to the other tissues containing this enzyme.					
14. SUBJECT TERMS <div style="text-align: right;">92-23379 418814 5p.</div>					
15. NUMBER OF PAGES				16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED		18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED		19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	
20. LIMITATION OF ABSTRACT UNLIMITED					

\*\*\* DTIC DATA \*\*\*

P. R. NUMBER: FQ8671-8901725  
 PROPOSAL NUMBER: 88NL262  
 TYPE SUBMISSION: FINAL Doc# 8262e, page 15  
 INST. CONTROL NUMBER: AFOSR-89-0190  
 INSTITUTION: American College of Toxicology  
 P.I. NAME: Ms Alexandra Ventura  
 INVENTION IND: NONE  
 PROJECT/TASK: 2312 A5  
 PROGRAM MANAGER: DR JIMMY CORNETTE

\*\*\*\*\*

3. Progress:

@25@UNARR. 502270 AFOSR-89-0190 PROG FROM 01 Dec 88 TO 31 May 90.

Organ specific distribution of carboxylesterases (Western blotting) was determined to be liver ] lung = testes -= fat ] pancreas ] kidney. Carboxylesterase distribution among cell types of the testes was examined by in situ hybridization techniques. Results were inconclusive, as both the probe and the control hybridized to tissue macromolecules. More refinement of this technique should provide better results. Other accomplishments include examination of the down-regulation of carboxylesterase levels by glucocorticoids. Apparently esterase levels are most dramatically down-regulated (approximately 6-fold) by dexamethasone phosphate (60 mg/kg x 5 days, i.p.) in the testes compared to the other tissues containing this enzyme.

Accession For	
NTIS	CRA&I
DTIC	TAB
Unannounced	
Justification	
By	
Distribution/	
Availability Codes	
Dist	Availability Codes
A-1	

CC: Cerveny

file  
2312/A

Rochelle M. Long, Ph.D.  
Dept. of Pharm. and Tox.  
School of Pharmacy  
University of Maryland

SETAC  
NI (AL)

# PROGRESS REPORT

8-1-89 TO 7-31-90


Air Force Office of Scientific Research

New Investigator Award

"Carboxylesterases of the Testes: Role in Activation of Toxicants"

## Progress towards proposed Specific Aims:

1. Organ specific distribution of the carboxylesterases was determined by Western blotting using anti-carboxylesterase antibodies and homogenates (20 ug total protein) of the various tissues. Relative abundance of the carboxylesterases was determined to be liver > lung = testes = fat > pancreas > kidney (see Fig. 1). At least 2 distinct bands were visualized in all of these tissues except the kidney, which is suggestive of multiple carboxylesterases forms, all of which are immunoreactive with the polyclonal antibodies.




2. Carboxylesterase distribution among cell types of the testes was examined by in situ hybridization techniques. Tissue slices were prepared as paraformaldehyde or immunobed sections. Samples were pre-hybridized and then hybridized with a <sup>32</sup>P-labeled antisense RNA or a sense RNA (control) made from a carboxylesterase clone inserted into the vector pGEM, in order to detect carboxylesterase-related sequences. Initial results were inconclusive, as both the probe and the control hybridized to tissue macromolecules (see Fig. 2). More refinement of this technique (to establish increased stringency without releasing the tissue section from the glass slide) should provide better results.

antisense      sense



3. Establishment of the co-cultures of testicular cell types was deferred, due to down time while the lab at University of Maryland was being set up and staffed. This method will be re-established in Dr. Robert Chapin's laboratory at NIEHS and the questions with DEHP and TOCP will be addressed collaboratively. These studies are pending resolution of the carboxylesterase localization studies, described in 2. above.

4. Other accomplishments related to the specific aims proposed in this grant include examination of the down-regulation of carboxylesterase levels by glucocorticoids (funded by the PMAF grant, identified below). Apparently esterase levels are most dramatically down-regulated (approximately 6-fold) by dexamethasone phosphate (60 mg/kg x 5 days, i.p.) in the testes (see Fig. 3) compared to the other tissues containing this enzyme.



#### Personnel:

Staffing for this project included Ms. Maria Calabrese (undergraduate student, part-time 1 day per week for the year), Mr. Jian-Ming Mei (graduate student, part-time for 6 weeks), and Mr. Tom Cooney (prospective graduate student, part-time for 4 weeks). No full-time staff was available during the first project year.

#### Budget Report:

The purchases of equipment and supplies to furnish the laboratory were in accordance with the modified budget submitted to the Society of Toxicology Office and approved by Joan Cassedy and Jan Cervany (copy attached). The final budget report will be available after the conclusion of the grant period, and can be obtained from Michael Gentry, Dept. of Pharmacology and Toxicology, School of Pharmacy, 20 N. Pine St., University of Maryland, Baltimore MD 21201, (301) 328-2976.

#### Manuscripts and Abstracts:

1. "Comparisons Between Rat and Human Liver Carboxylesterases", presentation at the 1990 IUPHAR Meeting, Amsterdam, The Netherlands.

2. "Human Liver Carboxylesterase: cDNA Cloning and Sequencing", presentation at the 1990 FASEB Meeting, Washington, DC.

3. "Human Liver Carboxylesterase: cDNA Cloning and Sequencing and Evidence for a Multigene Family", manuscript in preparation for submission to FEBS Letters.

Other Support:

"Distribution and Regulation of Carboxylesterases", Starter Grant from the Pharmaceutical Manufacturers Association Foundation, \$10,000/1 year (starting 1-1-90).

Extensions:

This project will not be continued after the conclusion of the grant period. The new address for the principal investigator will be (effective 8-1-90):

Dr. Rochelle M. Long  
NIGMS, NIH  
Westwood Bldg. Rm. 919  
Bethesda, MD 20892  
(301) 496-7707